



UNIVERSITI PUTRA MALAYSIA

**BIO-EFFICACY OF COMPOST EXTRACTS FOR THE CONTROL OF
CHOANEPHORA WET ROT OKRA (ABELMOSCHUS ESCULENTUS
L.)**

YASMEEN SIDDIQUI.

FP 2005 13

**BIO-EFFICACY OF COMPOST EXTRACTS FOR THE CONTROL OF
CHOANEPHORA WET ROT OF OKRA (*Abelmoschus esculentus* L.)**

By

YASMEEN SIDDIQUI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the
Degree of Doctor of Philosophy**

September 2005



DEDICATION

**To my beloved parents and husband without their support and inspiration
this goal could have not been achieved.**



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Doctor of Philosophy

**BIO-EFFICACY OF COMPOST EXTRACTS FOR THE CONTROL OF
CHOANEPHORA WET ROT OF OKRA (*Abelmoschus esculentus* L.)**

By

YASMEEN SIDDIQUI

September 2005

Chairman: Professor Sariah Meon, PhD

Faculty : Agriculture

Bio-efficacy of aqueous extracts prepared from rice straw (RST) and empty fruit bunch of oil palm (EFB) composts for the control of Choanephora wet rot of okra (*Abelmoschus esculentus* L.) was studied. The physiochemical and microbiological properties of both the finished composts showed that they were within the range of compost standards. RST compost extracts has higher N, P, K content of 1.14, 3.36, and 1.42 % with respect to EFB extract with 1.01, 2.51, and 1.20 %, respectively. The aqueous extracts produced therefore are suitable as foliar sprays for the improvement of growth of okra plants. Twelve days of brewing was the best extraction time based on the properties of the compost extracts obtained and their ability to inhibit the growth of *C. cucurbitarum* *in vitro*. The percent germination was reduced by 80 to 83 % in the presence of RST and EFB extracts. The effect was further demonstrated by using filter (0.2 µm) and heat - sterilized extracts (autoclaving at 121°C for 20 minutes) for assessing their potential effect on mycelial growth, conidial germination of *C. cucurbitarum*



and production of inducible resistance related compounds in pre – treated plants. Non – sterilized extracts from both composts gave significantly reduced ($P \leq 0.05$) mycelial growth of *C. cucurbitarum* (84 and 79 % for RST and EFB extracts, respectively) and enhanced host resistance through production of inducible compounds such as total phenols, peroxidase (PO), polyphenol oxidase (PPO) and phenylalanine ammonia lyase (PAL) in okra plants. The efficacy was perceptibly reduced by filter sterilizing the RST and EFB extracts to 57 and 55 %, respectively, whereas heat sterilization nullifies their activity totally, suggesting that the activity of both extracts were biological in nature. The higher activity of filter-sterilized extracts associated with mycelial growth inhibition and percentage disease reduction as compared to heat – sterilized extracts infers that chemical factors and nutrient composition of the extracts also played a role in the activity. Humic acids (active component of humic substances in compost) extracted from RST and EFB compost extracts when tested *in vitro* reduced mycelial growth of *C. cucurbitarum* by 100 and 64.2 % at 1% concentration, respectively. The bio-efficacy was further enhanced by fortifying the extracts with *T. harzianum*, an established biocontrol agent against many pathogens. *Trichoderma* - fortified extracts (3×10^{-7} cfu ml⁻¹) from RST when tested on okra in glass house and field conditions showed significant increase in plant growth and yield. There was reduction of 91.9 % in Choanephora wet rot of okra at 12 weeks of assessment in field and was comparable to Dithane M- 45® and significantly higher to *T. harzianum* fortified EFB extracts.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KEBERKESANAN BIOLOGI EKSTRAK KOMPOS UNTUK PENGAWALAN
REPUT BASAH CHOANEPHORA BENDI (*Abelmoschus esculentus*. L)**

Oleh

YASMEEN SIDDIQUI

September 2005

Pegerusi : Profesor Sariah Meon, PhD

Fakulti : Pertanian

Keberkesanan biologi ekstrak akueus yang disediakan daripada kompos jerami padi (RST) dan tandan kelapa sawit (EFB) untuk mengawal reput basah *Choanephora* pada pokok bendi (*Abelmoschus esculentus* L.) telah dikaji. Ciri-ciri kimia-fizik dan mikrobiologi yang terdapat pada kedua-dua kompos ini menepati julat piawai kompos. Ekstrak kompos daripada RST mempunyai kandungan N,P,K yang tinggi, masing-masing pada 1.14, 3.36 dan 1.42% berbanding dengan ekstrak kompos EFB iaitu 1.01, 2.51 dan 1.20 %. Oleh itu, ekstrak akueus yang dihasilkan sesuai sebagai semburan daun untuk memperbaiki pertumbuhan pokok bendi. Fermentasi selama 12 hari merupakan tempoh pengekstraktan yang paling baik berdasarkan ciri-ciri ekstrak yang dihasilkan dan keupayaannya untuk merencat pertumbuhan *C. cucurbitarum* secara *in vitro*. Peratus percambahan telah menurun dari 80 kepada 83 % dengan kehadiran ekstrak RST dan EFB. Kesan seterusnya

dibuktikan melalui penggunaan ekstrak yang disterilkan dengan turasan ($0.02\mu\text{m}$) dan haba (autoklaf pada 121°C selama 20 minit) untuk menilai potensi keberkesanannya terhadap pertumbuhan miselium, percambahan konidia *C. cucurbitarum*, dan penghasilan kompaun aruhan keresistanan pada pokok bendi yang dirawat dengan ekstrak kompos. Ekstrak kedua-dua kompos yang tidak disterilkan memberi keputusan yang signifikan ($P \leq 0.05$) merujuk kepada pengurangan pertumbuhan miselium *C. cucurbitarum* (masing-masing, 84 dan 79% untuk ekstrak RST dan EFB) dan peningkatan ketahanan perumah melalui stimulasi sebatian aruhan seperti fenol, peroksidase (PO), polifenol oksidase (PPO) dan fenilalanine ammonia liase (PAL) pada pokok bendi. Pensterilan melalui turasan mengurangkan keberkesanan ekstrak manakala pensterilan haba telah merencat keseluruhan keaktifan biologinya, mencadangkan aktiviti ekstrak mungkin dipengaruhi oleh ciri biologi semulajadi. Manakala Keberkesanan yang tinggi pada aktiviti ekstrak yang disterilkan melalui turasan berkaitan perencatan pertumbuhan miselium dan peratus pengurangan penyakit jika dibandingkan dengan ekstrak yang disterilkan dengan haba, menunjukkan kemungkinan peranan faktor kimia dan kandungan nutrien dalam aktiviti biologi ekstrak. Asid humik (komponen aktif bahan humik dalam kompos) yang diekstrak daripada kompos RST dan EFB apabila diuji secara *in vitro* telah merencat pertumbuhan miselium *C. cucurbitarum* kepada 100 dan 64% pada 1% kepekatan. Keberkesanan biologi diperkukuhkan seterusnya melalui penambahan *T. harzianum* agen, biologi yang telah teruji keberkesanannya terhadap kebanyakan patogen kulat. Ekstrak RST yang diperkukuhkan dengan *Trichoderma*, ($3 \times 10^{-7}\text{cfu ml}^{-1}$) apabila diuji pada pokok bendi di rumah kaca

dan ladang memberi peningkatan yang signifikan dalam pertumbuhan pokok dan hasil, dan mengurang/kan 91.9% reput basah Choanephora pada minggu ke-12 penilaian jika dibandingkan dengan Dithane M-45® dan peningkatan yang signifikan dari ekstrak EFB yang diperkukuhkan dengan *T. harzianum*.

ACKNOWLEDGEMENTS

My sincere gratitude goes to Professor Dr. Sariah Meon, the chairman of my supervisory committee for her dedicated efforts and scholarly guidance during the entire period of research and preparation of thesis. Her countless patience, encouragement, and generosity are commendable.

I am also grateful to Professor Dr. Mawardi Rahmani and Associate Professor Dr. Razi Ismail, the members of supervisory committee for their assistance at the various stages of my research.

I am thankful to all the staff in the pathology and chemistry laboratories, for their technical assistance.

I owe the members of my family particularly my parents, husband, my little master Hamzah and my father-in-law an incalculable debt for their love, care, sacrifices, spiritual support through their prayers and patience during the conduct of this research.

Best of all, all praises and thanks be to Allah the most beneficent and merciful, the creator of this magnificent universe and its constituents for the immeasurable guidance, good health, and direction. His name shall ever be praised and glorified.




I certify that an Examination Committee met on 8th September 2005 to conduct the final examination of Yasmeen Siddiqui on her Doctor of Philosophy thesis entitled "Bio-Efficacy of Compost Extracts for the Control of Choanephora Wet Rot of Okra (*Abelmoschus esculentus* L.)" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Kamaruzaman Sijam, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Halimi Mohd Saud, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Jugah Kadir, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Ibrahim Che Omar, PhD
Professor
School of Biological Sciences
Universiti Sains Malaysia
(External Examiner)



ZAKARIAH ABD RASHID, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **27 DEC 2005**

This thesis submitted to the Senate of Universiti Pertanian Malaysia and has been accepted as fulfillment of the requirements for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follow:

SARIAH MEON, PhD

Professor

Faculty of Agriculture

Universiti Putra Malaysia.

(Chairman)

Mawardi Rahmani, PhD

Professor

Faculty of Science

Universiti Putra Malaysia.

(Member)


Razi Ismail, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Member)



AINI IDERIS, PhD

Professor / Dean


School of Graduate Studies

Universiti Putra Malaysia

Date: 12 JAN 2006

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.


YASMEEN SIDDIQUI

Date: 6/11/05.

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xvii
LIST OF FIGURES	xix
LIST OF PLATES	xxii
LIST OF ABBREVIATIONS	xxiv

Chapter

1	INTRODUCTION	1.1
1.1	Hypothesis	1.6
1.2	Research Objectives	1.7
2	LITERATURE REVIEW	2.1
2.1	Shoot and Flower Blight	2.1
2.2	Choanephora Wet Rot	2.2
2.2.1	Symptoms	2.3
2.2.2	Host Range and Epidemiology	2.5
2.2.3	Control	2.6
2.3	Agro-wastes as Raw Materials for Composting	2.7
2.3.1	Oil Palm Wastes	2.8
2.3.2	Paddy Wastes	2.10
2.3.3	Chicken Manure	2.10
2.4	Composting Process	2.11
2.4.1	Bioconversion of Organic matter	2.14
2.4.2	Factors Affecting Composting Process	2.18
2.4.2.1	Particle size of materials	2.18
2.4.2.2	Moisture	2.19
2.4.2.3	Aeration	2.20
2.4.2.4	Temperature	2.21
2.4.2.5	pH	2.23
2.4.2.6	Carbon-Nitrogen Ratio	2.23
2.5	Compost Maturity	2.25
2.6	Compost Extracts	2.27
2.6.1	Composition of compost extracts	2.28
2.6.1.1	Beneficial Microorganisms	2.28
2.6.1.2	Humic Substances	2.30
2.7	Beneficial Effects of Compost and Compost	



	Extracts	2.34
	2.7.1 Improvement of plant growth	2.34
	2.7.2 Disease Suppression	2.36
	2.7.3 Mechanism of disease suppression	2.41
2.8	Improvement of Compost Extracts	2.48
3	EFFECTS OF PHYSIOCHEMICAL PROPERTIES OF RST AND EFB ON QUALITY OF COMPOST AND COMPOST EXTRACTS	
3.1	Introduction	3.1
3.2	Materials and Methods	3.3
3.2.1	Preparation of Substrates for Composting	3.3
	Grinding and Shredding	
3.2.2	Physiochemical Analyses of Composting	
	Materials	3.4
	Bulk Density	3.4
	Water Retention	3.5
	Particle Size	3.6
	Moisture Content	3.6
	pH and Electrical Conductivity	3.6
	Total Carbon	3.7
	Total Nitrogen	3.7
	C / N ratio	3.8
3.2.3	Composting Process	3.8
3.2.3.1	Temperature and Oxygen level	3.10
3.2.3.2	Nutrient Content and Heavy metal Accumulation	3.11
3.2.3.3	Microbial Population	3.12
3.2.4	Determination of Maturity of Compost	3.12
3.2.4.1	Evaluation of Toxicity of Compost Extracts	3.13
3.2.4.2	Effect of Compost on Seed Germination and Growth Performance of Okra	3.14
	Shoot and Root Length	3.15
	Dry Weight of Shoot and Root	3.15
	Number of Leaves	3.15
	Total Leaf Area	3.15
3.2.5	Statistical Analyses	3.16
3.3	Results and Discussion	3.17
3.3.1	Physiochemical Properties of Composting	
	Materials	3.17
	Structure and Texture	3.17
3.3.2	Composting Process	3.20
3.3.2.1	Colour Changes and Odour	3.20
3.3.2.2	Particle Size Distribution	3.22

3.3.2.3	Bulk Density	3.22
3.3.2.4	Water Retention	3.23
3.3.2.5	Aeration and Temperature	3.24
3.3.2.6	pH and Electrical Conductivity	3.28
3.3.2.7	Moisture Content	3.31
3.3.2.8	Total Carbon, Nitrogen and C/N ratio	3.32
3.3.2.9	Accumulation of Nutrients and Heavy Metals	3.34
3.3.3	Microbial Population in Finished Compost	3.36
3.3.4	Maturity Bioassay	3.39
3.3.4.1	Seed Germination	3.39
3.3.4.2	Seed Germination and Plant Growth in Compost	3.41
4	THE MECHANISMS OF SUPPRESSION OF <i>Choanephora cucurbitarum</i> BY RST AND EFB COMPOST EXTRACTS	
4.1	Introduction	4.1
4.2	Materials and Methods	4.3
4.2.1	Preparation of Compost Extracts	4.3
4.2.1.1	pH, Electrical Conductivity, Nutrients and Heavy metals Accumulation in Compost Extracts.	4.4
4.2.1.2	Microbial Population in Compost Extracts	4.4
4.2.2	Effect of RST and EFB Compost Extracts on <i>C. cucurbitarum</i>	4.5
4.2.2.1	Isolation of Pathogen and Preparation of Conidial Suspension	4.5
4.2.2.2	Determination of Optimal Time for Extraction	4.5
4.2.3	Effects of RST and EFB Compost Extracts on Mycelial Growth and Conidial Germination of <i>C. cucurbitarum</i>	4.6
4.2.3.1	Conidial Germination of <i>C. cucurbitarum</i>	4.6
4.2.3.2	Mycelial Growth	4.7
	Poison Food Method	4.7
	Bilayer Disc Method	4.8
4.2.4	Effects of Compost Extracts on Production of Inducible Compounds in Okra as Indicator of Resistance Mechanism	4.9
4.2.4.1	Effect of Compost Extracts on Disease Development	4.11
4.2.4.2	Determination of Total Phenolic Compounds	4.12
4.2.4.3	Determination of Enzymatic Activities	4.12
	Peroxidase Assay (PO)	4.13

	Polyphenol Oxidase Assay (PPO)	4.14
	Phenylalanine Ammonia Lyase (PAL)	4.14
4.3	Results and Discussion	4.16
4.3.1	Nutrients and Heavy metals, pH and Electrical Conductivity	4.16
4.3.2	Microbial Composition of RST and EFB Compost Extracts	4.19
4.3.3	Effect of Extraction Time on Conidial Germination	4.22
4.3.4	Effect of Compost Extracts on Conidial Germination and Mycelial growth of <i>C. cucurbitarum</i>	4.23
4.3.5	Effect of Compost Extracts on Disease Incidence and Production of Inducible Compounds	4.32
4.3.5.1	Disease Development	4.32
4.3.5.2	Production of Inducible Compounds as an Indicator of Disease Resistance in Okra Plants	4.36
5	BIOLOGICAL ACTIVITY OF HUMIC ACIDS FROM RST AND EFB COMPOST EXTRACTS	
5.1	Introduction	5.1
5.2	Materials and Methods	5.3
5.2.1	Extraction and Purification of Humic Acids	5.3
5.2.2	Fractionation of Humic Acids	5.4
5.2.3	Spectral Analysis	5.5
5.2.4	Biological Activity of Humic Acids Tested on <i>C. Cucurbitarum</i>	5.5
5.3	Results and Discussion	5.7
5.3.1	Extraction of Humic Acids	5.7
5.3.2	Humic Acid Fractions	5.8
5.3.3	Spectral Characteristics of Humic Acids	5.9
5.3.4	Biological Activity of Humic Acids on Mycelial Growth of <i>C. cucurbitarum</i>	5.11
6	EFFECT OF <i>TRICHODERMA</i> – FORTIFIED RST AND EFB COMPOST EXTRACTS ON PLANT GROWTH AND CONTROL OF CHOANEPHORA WET ROT OF OKRA	
6.1	Introduction	6.1
6.2	Materials and Methods	6.3
6.2.1	Isolation and Identification	6.3
6.2.2	Screening for Antagonistic Activity of <i>Trichoderma</i> Isolates against <i>C. cucurbitarum</i>	6.3
6.2.3	Effect of <i>Trichoderma</i> - Fortified Compost Extracts	

	on Growth Performance of Okra and Development of Choanephora Wet Rot	6.4
	Plant Materials and Growing Medium	6.4
	6.2.3.1 Growth Responses	6.6
	6.2.3.2 Disease Incidence	6.6
	6.2.3.3 Physiological Responses	6.6
6.2.4	Effect of <i>Trichoderma</i> -Fortified Compost Extracts On Yield and Disease Severity	6.8
	Field Preparation and Planting Material	6.9
	6.2.4.1 Disease Severity	6.11
	6.2.4.2 Effect of Compost Extracts on Yield and Yield Components of Okra	6.12
6.3	Results and Discussion	6.14
6.3.1	Isolation and Identification of <i>Trichoderma</i> from RST and EFB Compost	6.14
6.3.2	<i>In vitro</i> Screening for <i>Trichoderma</i> Isolates Effective against <i>C. cucurbitarum</i>	6.14
6.3.3	Effect of <i>Trichoderma</i> - Fortified Compost Extracts on Growth Performance of Okra and Development of Choanephora Wet Rot	6.16
	6.3.3.1 Growth Performance	6.16
	6.3.3.2 Disease Incidence	6.21
	6.3.3.3 Physiological Responses	6.25
6.3.4	Effect of <i>Trichoderma</i> - Fortified Compost Extracts on Yield and Disease Severity	6.30
	6.3.4.1 Disease Severity	6.30
	6.3.4.2 Effect of <i>Trichoderma</i> - fortified Compost Extracts on Yield and Yield Components of Okra	6.37
7	GENERAL DISCUSSION AND CONCLUSIONS	7.1
	REFERENCES	R.1
	APPENDICES	A.1
	BIODATA OF THE AUTHOR	B.1

LIST OF TABLES

Table	Page
2.1. Effects of compost on growth and yield of selected vegetable crops.	2.35
3.1. Compost to peat combinations of plant growing media in container mix.	3.14
3.2. Physiochemical properties of grounded Rice Straw (RST) and Empty Fruit Bunch (EFB) of oil palm before composting Process.	3.19
3.3. Physiochemical properties of chicken manure	3.19
3.4. Some physiochemical properties of composts produced from RST and EFB supplemented with chicken manure.	3.24
3.5 Accumulation of nutrients and heavy metals throughout the composting process of RST and EFB supplemented with chicken manure.	3.36
3.6 Microbial Population in finished RST and EFB Compost supplemented with chicken Manure.	3.37
3.7 Vegetative growth of okra seedlings sown in different combinations of compost and peat 4 weeks after sowing.	3.44
4.1 Treatments for the evaluation of compost extracts for the production of inducible compounds and disease incidence in okra plants.	4.9
4.2 Nutrients and heavy metals content in RST and EFB compost extracts for the different extraction time.	4.17
4.3 Disease incidence, epidemic rates and area under disease progress curve (AUDPC) for <i>Choanephora</i> wet rot treated with compost extracts (different methods of sterilization) after 14 days of inoculation with <i>C. cucurbitarum</i>	4.36
5.1 Recovery of humic acid extracted from RST and EFB compost extracts using C 18 column with 100 Å pore size. Analytical grade sodium salt of humic acid Na HA was used as standard.	5.9

5.2	Absorption bands in the FTIR spectra of humic acids derived from RST and EFB using KBr pellets. The analytical grade humic acid salt NaHA was used as standard.	5.11
6.1	Treatments for evaluating the effectiveness of <i>Trichoderma</i> - fortified compost extracts on growth performance and <i>Choanephora</i> wet rot of okra.	6.5
6.2	Physiochemical characteristics of Serdang series soil from field No.10, Universiti Putra Malaysia.	6.8
6.3	Effect of different compost extracts on root length, shoot length, number of leaves and leaf area of <i>Choanephora</i> inoculated and uninoculated okra plants on weekly basis.	6.19
6.4	Disease progress rates and area under disease progress curve (AUDPC) for <i>Choanephora</i> wet rot on okra treated with RST and EFB compost extracts at 4 weeks after inoculation.	6.23
6.5	Disease progress rates and area under disease progress curve (AUDPC) for <i>Choanephora</i> wet rot on okra treated with different compositions of compost extracts at 12 weeks of assessment.	6.33
6.6	Effect of foliar application of different compost extracts on number of buds, flowers, pods, pod length and weight of pods at weekly intervals.	6.40
6.7	Mean values of diseased pods, total yield, marketable yield and percent loss in yields of okra plants treated with different compost extracts after 12 weeks of assessment.	6.41

LIST OF FIGURES

Figure	Page
3.1 Variation in level of oxygen throughout the composting process of rice straw (RST) and empty fruit bunch (EFB) of oil palm supplemented with chicken manure.	3.26
3.2. Variation in temperature throughout the composting process of rice straw (RST) and empty fruit bunch (EFB) of oil palm supplemented with chicken manure.	3.28
3.3. Variation in pH during composting of rice straw (RST) and empty fruit bunch (EFB) of oil palm supplemented with chicken manure.	3.29
3.4 Variation in electrical conductivity (EC) throughout composting process of rice straw (RST) and empty fruit bunch (EFB) of oil palm supplemented with chicken manure.	3.30
3.5 Variation in moisture content throughout the composting process of rice straw (RST) and empty fruit bunch (EFB) of oil palm supplemented with chicken manure.	3.32
3.6 Variation in total carbon, total nitrogen and C/N ratio throughout the composting process of rice straw (RST) and empty fruit bunch (EFB) of oil palm supplemented with chicken manure.	3.34
3.7 Percentage germination of chinese cabbage seeds treated with rice straw (RST) and empty fruit bunch (EFB) of oil palm compost extracts.	3.40
3.8 Percentage germination of okra seeds treated with rice straw (RST) and empty fruit bunch (EFB) of oil palm compost extracts.	3.40
3.9 Percentage germination of okra seeds sown in different combinations of rice straw (RST) and empty fruit bunch (EFB) compost with peat, seven days after sowing.	3.42
4.1a Variation in pH of RST (Rice straw) and EFB (Empty fruit bunch) compost extracts at different extraction times.	4.18

4.1 b	Variation in electrical conductivity (EC) of RST (Rice straw) and EFB (Empty fruit bunch) compost extracts at different extraction times.	4.19
4.2 A	(a) <i>Trichoderma</i> spp ($\times 10^5$) (b) <i>Aspergillus</i> spp ($\times 10^6$) isolated from rice straw (RST) and empty fruit bunch (EFB) compost extracts at different extraction times.	4.21
4.2 B	(a) <i>Penicillium</i> spp ($\times 10^4$) (b) Total Actinomycetes ($\times 10^3$) (c) Total Bacteria ($\times 10^{11}$) and (d) <i>Pseudomonas</i> ($\times 10^4$) in rice straw (RST) and empty fruit bunch (EFB) compost extracts at different extraction times.	4.22
4.3	Effect of extraction time of rice straw (RST) and empty fruit bunch (EFB) compost extracts on percentage inhibition of conidial germination of <i>C. cucurbitarum</i> at different extraction times.	4.23
4.4	Effect of sterilization techniques of rice straw compost (RST) and empty fruit bunch (EFB) compost extracts on inhibition of conidial germination of <i>C. cucurbitarum</i> .	4.25
4.5	Effect of sterilization techniques of rice straw (RST) and empty fruit bunch (EFB) compost extracts on percentage inhibition of mycelial growth of <i>C. cucurbitarum</i> .	4.27
4.6	Disease incidence (%) in okra plants <i>var five anchors</i> pre - treated with compost extracts and challenged with <i>C. cucurbitarum</i> .	4.35
4.7	Total phenolics in okra leaves pre-treated with compost extracts (A) pre - treated with compost extracts and challenged with <i>C. cucurbitarum</i> (B) in relation to disease incidence (C).	4.39
4.8	Total peroxidase (PO) activity in okra leaves pre - treated with compost extracts (A) pre - treated with compost extracts and challenged with <i>C. cucurbitarum</i> (B) in relation to disease incidence (C).	4.43

4.9	Polyphenol oxidase (PPO) activity in okra leaves pre - treated with compost extracts (A) pre - treated with compost extracts and challenged with <i>C. cucurbitarum</i> (B) in relation to disease incidence (C).	4.46
4.10	Phenylalanine Ammonia Lyase (PAL) activity in okra leaves pre - treated with compost extracts (A) pre - treated with compost extracts and challenged with <i>C. cucurbitarum</i> (B) in relation to disease incidence (C).	4.48
5.1	Effect of different concentrations of humic acids extracted from RST and EFB compost extracts on percentage inhibition in mycelial growth of <i>C. cucurbitarum</i> after five days of incubation at $28 \pm 2^{\circ}$ C.	5.12
6.1	Field layout for evaluating the effect of <i>Trichoderma</i> – fortified compost extracts on pod yield and disease severity.	6.10
6.2	<i>In vitro</i> antagonistic activity of different isolates of <i>Trichoderma harzianum</i> isolated from rice straw (RST) and empty fruit bunch (EFB) of oil palm composts against <i>C. cucurbitarum</i> .	6.16
6.3	Disease incidence in okra plants inoculated with <i>C.cucurbitarum</i> .	6.22
6.4	Net photosynthesis rate ($\text{m mol m}^{-2}\text{s}^{-1}$) of okra plants inoculated with <i>C. cucurbitarum</i> (B) and uninoculated (C) in relation to disease incidence (A).	6.27
6.5	Total chlorophyll content (mg cm^{-2}) in (A) Inoculated and (B) Uninoculated leaves of okra plants treated with different compost extracts.	6.29
6.6	Disease severity assessed in okra plants caused by <i>Choanephora</i> wet rot on okra plants treated with different compost extracts at weekly interval.	6.31
6.7	Correlation between disease severity and total yield of okra plants at 12 weeks of harvesting.	6.42

LIST OF PLATES

Plates	Page
3.1 Composting bins supplied with aeration system using three perforated PVC tubes connected to an air compressor.	3.9
3.2 Temperature and oxygen indicators placed in the composting bins.	3.11
3.3 Composting materials: rice straw (RST) and empty fruit bunch (EFB) of oil palm after grinding process	3.17
3.4 Finished compost after 120 Days of composting Process (A) Rice straw (RST) and, (B) Empty Fruit Bunch of Oil Palm (EFB)	3.21
3.5 Vegetative growth of okra grown in different compositions of rice straw (RST) compost and peat, 4 weeks after sowing.	3.45
3.6 Vegetative growth of okra grown in different compositions of empty fruit bunch (EFB) of oil palm compost and peat, 4 weeks after sowing.	3.45
4.1 Compost extracts prepared from three-month-old (A) Rice Straw (RST) and (B) Empty Fruit Bunch (EFB) compost.	4.3
4.2 Effect of RST compost extracts on conidial germination of <i>C. cucurbitarum</i> after 24 h of incubation (a) non-sterilized, (b) filter-sterilized, (c) heat-sterilized compost extracts and (d) sterilized distilled water, mg- malformed germ tube; g- germ tube.	4.26
4.3 Effect of sterilization techniques of rice straw (RST) and empty fruit bunch (EFB) compost extracts on mycelial growth of <i>C. cucurbitarum</i> , 5 days after incubation. A) Non-sterilized, B) Filter-sterilized, C) Heat-Sterilized and D) Control (PDA).	4.28
4.4 Formation of clearing zones of inhibition in mycelial growth of <i>C. cucurbitarum</i> in the presence of A) non-sterilized, B) filter-sterilized C) heat-sterilized and D) Distilled water (Control).	4.30
4.5 Effect of RST compost extracts on the vegetative structure of <i>C. cucurbitarum</i> . a) Non- sterilized, b) Filter- sterilized and c) water (control).	4.31

4.6	Development of foliar-associated symptoms produced on okra leaves pre-treated with compost extracts and challenged with <i>C. cucurbitarum</i> .	4.34
5.1	Effect of different concentrations of humic acids extracted from RST and EFB compost extracts on percentage inhibition in mycelial growth of <i>C. cucurbitarum</i> after five days of incubation at $28 \pm 2^{\circ}\text{C}$.	5.13
6.1	<i>In vitro</i> antagonistic activity <i>Trichoderma</i> isolates isolated from rice straw (RST) and empty fruit bunch (EFB) of oil palm composts against <i>C. cucurbitarum</i> based on dual culture test.	6.15
6.2	Development of <i>Choanephora</i> wet rot caused by <i>C. cucurbitarum</i> after 4 weeks of assessment in okra plants receiving T2 (A), T4 (B), T7 (C), T5 (F) and T6 (E).	6.24
6.3	Compost extracts treated plants with a disease index score of 0 = healthy; 1 = 1 - 5% coverage by necrotic and chlorotic symptoms on lower mature leaves; 2 = 6 - 15 % coverage by necrotic and chlorotic symptoms on mature lower leaves extending towards stem; 3 = 16- 50 % area covered by water soaked lesions on young shoots and stem; 4 = 51 – 95% coverage by water soaked lesions and shiny mycelial growth on shoots, buds, flowers and fruits.	6.32
6.4	Okra plants infected with <i>Choanephora</i> wet rot showing infected shoot, bud and flower (A), severely infected okra plant in control (B), flower attached to pod after fruit set covered with a silvery, spine-like fungus with a dark purple pin heads (C) and healthy plant treated with <i>Trichoderma</i> -amended RST compost extract (D).	6.35

LIST OF ABBREVIATIONS

AgNO ₃	Silver Nitrate
ANOVA	Analysis of Variance
AUDPC	Area Under Disease Progress Curve
C	Carbon
Ca	Calcium
Cd	Cadmium
c.f.u	Colony Forming Units
CHB	Composted Hardwood Bark
CO ₂	Carbon dioxide
Cu	Copper
CRD	Completely Randomized Design
DI	Disease Incidence
EC	Electrical Conductivity
EFB	Empty Fruit Bunch
EPA	Environmental Protection Agency
EQA	Environmental Quality Act
Fe	Iron
Fe ³⁺	Ferric Ion
g cm ⁻²	Gram per Centimeter square
HCL	Hydrochloric acid
HF	Hydrogen Flouride
K	Potassium
kDa	Kilo Dalton
Kg ha ⁻¹	Kilogram per Hectare
LSD	Least Significant Difference
M	Molar
Mt	Metric Tonnes
Mt/ha	Metric tonnes per Hectare
mgL ⁻¹	Milligram per liter